

Adult ADHD Self-Report Scale (ASRS-v1.1) Symptom Checklist

Patient Name		Today's Date					
<p>Please answer the questions below, rating yourself on each of the criteria shown using the scale on the right side of the page. As you answer each question, place an X in the box that best describes how you have felt and conducted yourself over the past 6 months. Please give this completed checklist to your healthcare professional to discuss during today's appointment.</p>			Never	Rarely	Sometimes	Often	Very Often
1. How often do you have trouble wrapping up the final details of a project, once the challenging parts have been done?							
2. How often do you have difficulty getting things in order when you have to do a task that requires organization?							
3. How often do you have problems remembering appointments or obligations?							
4. When you have a task that requires a lot of thought, how often do you avoid or delay getting started?							
5. How often do you fidget or squirm with your hands or feet when you have to sit down for a long time?							
6. How often do you feel overly active and compelled to do things, like you were driven by a motor?							
Part A							
7. How often do you make careless mistakes when you have to work on a boring or difficult project?							
8. How often do you have difficulty keeping your attention when you are doing boring or repetitive work?							
9. How often do you have difficulty concentrating on what people say to you, even when they are speaking to you directly?							
10. How often do you misplace or have difficulty finding things at home or at work?							
11. How often are you distracted by activity or noise around you?							
12. How often do you leave your seat in meetings or other situations in which you are expected to remain seated?							
13. How often do you feel restless or fidgety?							
14. How often do you have difficulty unwinding and relaxing when you have time to yourself?							
15. How often do you find yourself talking too much when you are in social situations?							
16. When you're in a conversation, how often do you find yourself finishing the sentences of the people you are talking to, before they can finish them themselves?							
17. How often do you have difficulty waiting your turn in situations when turn taking is required?							
18. How often do you interrupt others when they are busy?							
Part B							

Patient Name: _____ DOB: _____ Date: _____

Beck's Depression Inventory

This depression inventory can be self-scored. The scoring scale is at the end of the questionnaire.

1.
 - 0 I do not feel sad.
 - 1 I feel sad.
 - 2 I am sad all the time and I can't snap out of it.
 - 3 I am so sad and unhappy that I can't stand it.
2.
 - 0 I am not particularly discouraged about the future.
 - 1 I feel discouraged about the future.
 - 2 I feel I have nothing to look forward to.
 - 3 I feel the future is hopeless and that things cannot improve.
3.
 - 0 I do not feel like a failure.
 - 1 I feel I have failed more than the average person.
 - 2 As I look back on my life, all I can see is a lot of failures.
 - 3 I feel I am a complete failure as a person.
4.
 - 0 I get as much satisfaction out of things as I used to.
 - 1 I don't enjoy things the way I used to.
 - 2 I don't get real satisfaction out of anything anymore.
 - 3 I am dissatisfied or bored with everything.
5.
 - 0 I don't feel particularly guilty.
 - 1 I feel guilty a good part of the time.
 - 2 I feel quite guilty most of the time.
 - 3 I feel guilty all of the time.
6.
 - 0 I don't feel I am being punished.
 - 1 I feel I may be punished.
 - 2 I expect to be punished.
 - 3 I feel I am being punished.
7.
 - 0 I don't feel disappointed in myself.
 - 1 I am disappointed in myself.
 - 2 I am disgusted with myself.
 - 3 I hate myself.
8.
 - 0 I don't feel I am any worse than anybody else.
 - 1 I am critical of myself for my weaknesses or mistakes.
 - 2 I blame myself all the time for my faults.
 - 3 I blame myself for everything bad that happens.
9.
 - 0 I don't have any thoughts of killing myself.
 - 1 I have thoughts of killing myself, but I would not carry them out.
 - 2 I would like to kill myself.
 - 3 I would kill myself if I had the chance.

20.

- 0 I am no more worried about my health than usual.
- 1 I am worried about physical problems like aches, pains, upset stomach or constipation
- 2 I am very worried about physical problems and it's hard to think of much else.
- 3 I am so worried about my physical problems that I cannot think of anything else.

21.

- 0 I have not noticed any recent change in my interest in sex.
- 1 I am less interested in sex than I used to be.
- 2 I have almost no interest in sex.
- 3 I have lost interest in sex completely.

Interpreting the Beck Depression Inventory

Now that you have completed the questionnaire, add up the score for each of the twenty-one questions by counting the number to the right of each question you marked. The highest possible total for the whole test would be sixty-three. This would mean you circled number three (3) on all twenty-one questions. Since the lowest possible score for each question is zero (0), the lowest possible score for the test would be zero. This would mean you circled zero (0) on each question. You can evaluate your depression according to the table below.

Total Score	Levels of Depression
1-10	There ups and downs are considered normal
11-16	Mild mood disturbance
17-20	Borderline clinical depression
21-30	Moderate depression
31-40	Severe depression
Over 40	Extreme depression

Patient Name: _____ DOB: _____ Date: _____

Beck Anxiety Inventory

Below is a list of common symptoms of anxiety. Please carefully read each item in the list. Indicate how much you have been bothered by that symptoms during the past month, including today, by circling the number in the corresponding space in the column next to each symptom.

	Not At All	Mildly but it didn't bother me much	Moderately – it wasn't pleasant at times	Severely – it bothered me a lot
Numbness or tingling	0	1	2	3
Feeling hot	0	1	2	3
Wobbliness in legs	0	1	2	3
Unable to relax	0	1	2	3
Fear of worst happening	0	1	2	3
Dizzy / lightheaded	0	1	2	3
Heart pounding/racing	0	1	2	3
Unsteady	0	1	2	3
Terrified or afraid	0	1	2	3
Nervous	0	1	2	3
Feeling of choking	0	1	2	3
Hands trembling	0	1	2	3
Shaky / unsteady	0	1	2	3
Fear of losing control	0	1	2	3
Difficulty in breathing	0	1	2	3
Fear of dying	0	1	2	3
Scared	0	1	2	3
Indigestion	0	1	2	3
Faint / lightheaded	0	1	2	3
Face flushed	0	1	2	3
Hot / Cold sweats	0	1	2	3
COLUMN SUMMARY				

Scoring – Sum each column, then sum column totals to achieve a grand score. Write that score here _____.

Interpretation

A grand sum between 0 – 21 indicates very low anxiety. That is usually a good thing. However, it is possible that you might be unrealistic in either your assessment which would be denial or that you have learned to “mask” the symptoms commonly associated with anxiety. Too little “anxiety” could indicate that you are detached from yourself, others or your environment.

A grand sum between 22 – 35 indicates moderate anxiety. Your body is trying to tell you something. Look for patterns as to when and why you experience the symptoms described above. For example, if it occurs prior to public speaking and your job requires a lot of presentations, you may want to find ways to calm yourself before speaking or let others do some of the presentations. You may have some conflict issues that need to be resolved. Clearly, it is not “panic” time but you want to find ways to manage the stress you feel.

A grand sum that exceeds 36 is a potential cause for concern. Again, look for patterns or times when you tend to feel the symptoms you have circled. Persistent and high anxiety is not a sign of personal weakness or failure. It is, however, something that needs to be proactively treated or there could be significant impacts to you mentally and physically. You may want to consult a counselor if the feelings persist.